




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 31/44, 31/36, 31/335, A01N 43/08, 43/30, 43/32, C07D 307/88, 317/46, 319/04, 319/08</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/62515</p> <p>(43) International Publication Date: 9 December 1999 (09.12.99)</p>
<p>(21) International Application Number: PCT/US99/11474</p> <p>(22) International Filing Date: 25 May 1999 (25.05.99)</p> <p>(30) Priority Data: 60/088,199 5 June 1998 (05.06.98) US</p> <p>(71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US).</p> <p>(72) Inventors: CHEN, Jie; 14 Martleshamheath Lane, Madison, CT 06443 (US). DEXTRAZE, Pierre; 6600 de la Bataille, Laprairie, Quebec J5R 3X8 (CA). DODIER, Marco; 320 Des Tulipes, Ste-Catherine, Quebec J0L 1E0 (CA). TAKAKI, Katherine, S.; 162 Acorn Drive, Middletown, CT 06457 (US).</p> <p>(74) Agent: ALGIERI, Aldo, A.; Bristol-Myers Squibb Company, 5 Research Parkway, P.O. Box 5100, Wallingford, CT 06492-7660 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: HETEROCYCLIC CIS CYCLOPROPANE DERIVATIVES AS MELATONERGIC AGENTS</p> <div style="text-align: center;">  <p>(I)</p> </div> <p>(57) Abstract</p> <p>There is provided a novel series of cis cyclopropane compounds of Formula (I), wherein R¹, R², R³, R⁴, X, Y, Z and m are as defined herein which bind to the human melatonin receptor and therefore are useful as melatonergic agents.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		


5 **HETEROCYCLIC CIS CYCLOPROPANE DERIVATIVES**
 AS MELATONERGIC AGENTS

10

BACKGROUND OF THE INVENTION

 The present invention pertains to novel substituted heterocyclic
15 cis cyclopropane derivatives having drug and bio-affecting properties
and to their preparation, pharmaceutical formulations and use. In
particular, the invention concerns benzodioxoles, benzofurans,
dihydrobenzofurans, dihydrobenzodioxanes and related derivatives
bearing aminoalkyl substituted cis cyclopropane groups. These
20 compounds possess melatonergic properties that should make them
useful in treating certain medical disorders.

 Melatonin (N-acetyl-5-methoxytryptamine) is a hormone which is
synthesized and secreted primarily by the pineal gland. Melatonin levels
25 show a cyclical, circadian pattern with highest levels occurring during the
dark period of a circadian light-dark cycle. Melatonin is involved in the
transduction of photoperiodic information and appears to modulate a
variety of neural and endocrine functions in vertebrates, including the
regulation of reproduction, body weight and metabolism in photoperiodic
30 mammals, the control of circadian rhythms and the modulation of retinal
physiology.



Melatonin

5 Recent evidence demonstrates that melatonin exerts its biological effects through specific receptors. Use of the biologically active, radiolabelled agonist [¹²⁵I]-2-iodomelatonin has led to the identification of high affinity melatonin receptors in the CNS of a variety of species. The sequences of two cloned human melatonin receptors have been
10 reported [Reppert, et al., Proc. Natl. Acad. Sci. 92, p. 8734-8738, (1995) and Reppert, et al., Neuron 13, p. 1177-1185, (1994)]. In mammalian brain, autoradiographic studies have localized the distribution of melatonin receptors to a few specific structures. Although there are significant differences in melatonin receptor distribution even between
15 closely related species, in general the highest binding site density occurs in discrete nuclei of the hypothalamus. In humans, specific [¹²⁵I]-2-iodomelatonin binding within the hypothalamus is completely localized to the suprachiasmatic nucleus, strongly suggesting the melatonin receptors are located within the human biological clock.


20

Exogenous melatonin administration has been found to synchronize circadian rhythms in rats (Cassone, et al., J. Biol. Rhythms, 1:219-229, 1986). In humans, administration of melatonin has been used to treat jet-lag related sleep disturbances, considered to be caused
25 by desynchronization of circadian rhythms (Arendt, et al., Br. Med. J. 292:1170, 1986). Further, the use of a single dose of melatonin to induce sleep in humans has been claimed by Wurtman in International Patent Application WO 94/07487, published on April 14, 1994.

30 Thus, melatonin agonists should be particularly useful for the treatment of sleep disorders and other chronobiological disorders. Melatonin agonists would also be useful for the further study of

melatonin receptor interactions as well as in the treatment of conditions
affected by melatonin activity, such as depression, jet-lag, work-shift
syndrome, sleep disorders, glaucoma, reproduction, cancer,
premenstrual syndrome, immune disorders, inflammatory articular
5 diseases and neuroendocrine disorders.


Aside from simple indole derivatives of melatonin itself, various
bicyclic structures have been prepared and their use as melatonin
ligands disclosed. In general these bicyclic amide structures can be
10 represented as:



wherein Z is an aryl or heteroaryl system attached by a two carbon
15 bridge to the amide group. Some specific examples follow.


Yous, et al. in European Patent Application EP-527,687A,
published on February 17, 1993, disclose as melatonin ligands
arylethylamines i,

20




wherein Ar' is, inter alia, a substituted or unsubstituted benzo[b]thiophen-
25 3-yl, benzimidazol-1-yl, benzo[b]furan-3-yl, 1,2-benzisoxazol-3-yl, 1,2-
benzothiazol-3-yl, or indazol-3-yl radical; R₁ is, inter alia, an alkyl or
cycloalkyl group; and R₂ is hydrogen or lower alkyl.

Yous, et al. in European Patent Application EP-506,539A,
30 published on September 30, 1992, claim ligands ii,




- wherein A is oxygen or sulfur; X is a methylene group or a bond; and R is
 5 H or lower alkyl when p is 1 and B is defined by the radical iii,



- 10 wherein R₁ is hydrogen or lower alkyl and R₂ is, inter alia, hydrogen, lower alkyl or cycloalkyl. Alternatively, R is defined by the radical iii when p is 0 or 1 and B is lower alkoxy.


- Several naphthalene derivatives have also been disclosed as
 15 melatonin ligands.

- Yous, et al. in European Patent Application EP-562,956A, published on September 29, 1993, disclose amide and urea naphthalene derivatives iv,
 20



- in which R is hydrogen or OR₄ wherein R₄ is, inter alia, hydrogen, alkyl, cycloalkyl, or cycloalkylalkyl; R₁ is hydrogen or COOR₅ wherein R₅ is
 25 hydrogen or alkyl; R₂ is hydrogen or alkyl; X is NH or a bond; and R₃ is, inter alia, alkyl, alkenyl, or cycloalkyl.

Langlois, et al., in Australian Patent Application AU-A-48729/93 disclose arylalkyl(thio)amides v as melatonergic ligands,



5

v


wherein R_1 is hydrogen or lower alkyl; R_2 is hydrogen, halogen, or lower alkyl; R_3 and R_4 are identical or different groups including, inter alia, hydrogen, halogen, or lower alkyl or R_3 and R_4 , together with the
 10 benzene ring which carries them, form a ring-system E_3 chosen from, inter alia, naphthalene, on the understanding that the portion of the ring-system E_3 formed by R_3 and R_4 and the two carbon atoms of the benzene ring which carry them is unhydrogenated or partially hydrogenated; R_5 is hydrogen or lower alkyl; and R_6 is,

15




wherein X is sulfur or oxygen and R_7 is, inter alia, lower alkyl or alkenyl.
 Compound vi is included as a specific example,

20


vi

Horn and Dubocovich in European Patent Application EP-420,064A, published on April 3, 1991, disclose 2-amidotetralins vii as melatonin ligands,

vii


wherein R₁ is, inter alia, hydrogen, lower alkyl, or lower alkoxy; R₂ is, inter alia, hydrogen, halogen, or lower alkoxy; R₃ is, inter alia, hydrogen, or lower alkyl; R₄ is, inter alia, lower alkyl, haloalkyl or cycloalkyl; and R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, lower alkyl or alkylaryl.

Lesieur et al., in EP-708,099A, published April 24, 1996, disclose compounds of structure viii, which are useful for the treatment of diseases caused by a melatonin imbalance.

viii


wherein --- is a single or double bond; R₁ = Me or MeNH; and X-Y = -CH(Me)-CH₂-, CH₂CH(OH)- or (CH₂)₃-.

North et al., in International Application WO 95/29173, published November 2, 1995, disclose naphthalene derivatives of structure ix:

ix


wherein R_1 is a group of the formula $CR_3R_4(CH_2)_pNR_5COR_6$; R_2 is
 5 hydrogen, halogen, C_{1-6} alkyl, OR_7 or CO_2R_7 ; and may be the same or
 different substituent when q is 2; R_3 , R_4 and R_5 , which may be the same
 or different, are hydrogen or C_{1-6} alkyl; R_6 is C_{1-6} alkyl or C_{3-7} cycloalkyl;
 R_7 is hydrogen or C_{1-6} alkyl; n is zero, 1 or 2; p is an integer of 1, 2, 3 or
 4; q is 1 or 2; and the dotted lines indicate the absence or presence of an
 10 additional bond. The North et al. compounds are taught to treat
 chronobiological disorders.

In International Application WO 95/17405, published on June 29,
 1995, North et al., disclose compounds of structure x and teach their use
 15 in the treatment of conditions related to the melatonin system.

x

20 wherein R_1 is hydrogen, halogen or C_{1-6} alkyl; R_2 is a group of formula
 $-CR_3R_4(CH_2)_pNR_5COR_6$; R_3 , R_4 and R_5 , which may be the same or
 different, are hydrogen or C_{1-6} alkyl; R_6 is C_{1-6} alkyl or C_{3-7} cycloalkyl; n
 is an integer of 2, 3 or 4; and p is an integer of 1, 2, 3 or 4.


Keavy, et al., in United States Patent No. 5,753,709, issued on May 19, 1998, disclose compounds of formula xi which are useful as melatonergic agents,



wherein X represents halogen, hydrogen, C₁₋₄alkyl or OR₅ wherein, inter alia, R₅ is hydrogen, C₁₋₂₀alkyl or C₄₋₂₀alkylcycloalkyl; Y represents
 10 hydrogen or halogen; X represents inter alia, hydrogen, halogen, cyano or aryl; R represents hydrogen, halogen or C₁₋₄alkyl or; R₁ represents hydrogen, C₁₋₄alkyl or benzyl and R₂ represents C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₆cycloalkyl, C₂₋₄alkoxyalkyl, C₁₋₄trifluoromethylalkyl or C₂₋₈alkylthioalkyl.

15

In International Application WO 97/43272, published on November 20, 1997, Ellis, et al., disclose compounds of structure xii as melatonin ligands.




wherein R¹ and R² represent hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl or aryl, R³ and R⁴ represent hydrogen, halogen or C₁₋₆alkyl or substituted aryl,
 25 R⁵ represents hydrogen or C₁₋₆alkyl, n is 0-2, m is 1-4 and the dotted line represents an additional bond.

The foregoing disclosures do not teach or suggest the novel melatonergic benzodioxole, benzofuran or dihydrobenzofurans of the present invention. The novel compounds of the present invention display melatonergic agonist activity.

5

SUMMARY OF THE INVENTION

The invention provides a novel series of cis cyclopropane
10 compounds of Formula I




wherein R¹, R², R³, R⁴, X, Y, Z and m are as defined below, including
15 hydrates and solvates thereof which bind to the human melatonergic receptor and therefore are useful as melatonergic agents in the treatment of sleep disorders, seasonal depression, shifts in circadian cycles, melancholia, stress, appetite regulation, benign prostatic hyperplasia and related conditions.

20

DETAILED DESCRIPTION OF THE INVENTION

The invention provides a novel series of cis cyclopropane
25 compounds of Formula I and solvates thereof having the formula:



wherein

- R¹ and R² each are independently hydrogen or halogen;
X is CH₂, CH or oxygen;
Y is CR⁵, CR⁵R⁶ or (CH₂)_n, with n = 1-2;
5 Z is CH₂, CH or oxygen;
m is 1 or 2;
R³ is hydrogen or C₁₋₄ alkyl;
R⁴ is C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₃ haloalkyl, C₂₋₆ alkenyl,
C₁₋₄ alkoxy(C₁₋₄)alkyl, C₁₋₄ alkylthio(C₁₋₄)alkyl or C₁₋₄
10 trifluoromethylalkyl; and
R⁵ and R⁶ each are independently hydrogen or C₁₋₄ alkyl.

The present invention also provides a method for the treatment of
sleep disorders and related conditions, which comprises administering a
15 therapeutically effective amount of a compound of Formula I or a solvate
or hydrate thereof.

R¹ and R² are selected from H and halogen (i.e., bromine,
chlorine, iodine or fluorine). It is most preferred that R¹ and R² be H.
20

X may be CH₂, CH (when a double bond is present) or oxygen.

Y is CR⁵ (when a double bond is present), CR⁵R⁶ or -(CH₂)_n- and
n is preferably 1 or 2.
25

Z may be CH₂, CH (when a double bond is present) or oxygen,
with oxygen being most preferred.

When X and Y are CH₂ and Z is oxygen or Z and Y are CH₂ and X
30 is oxygen, the compound is a dihydrobenzofuran. When X and Y are CH
and Z is oxygen or Z and Y are CH and X is oxygen, the compound is a
benzofuran. When X and Z are oxygen and Y is CH₂, the compound is a

benzodioxole. When X and Z are oxygen and Y is $(CH_2)_2$, the compound is a benzodioxane. Compounds in which X and Y are CH_2 and Z is oxygen are preferred.

5 m is 1 or 2, with m = 1 preferred.

R^4 is one of several types of groups. R^4 is selected from C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-3} haloalkyl, C_{2-6} alkenyl, C_{1-4} alkoxy(C_{1-4})alkyl, C_{1-4} trifluoromethylalkyl and C_{1-4} alkylthio(C_{1-4})alkyl groups. R^1 is
10 preferably C_{1-6} alkyl or C_{3-6} cycloalkyl.

R^3 is hydrogen or C_{1-4} alkyl. R^3 is preferably hydrogen.

R^5 and R^6 are hydrogen or C_{1-4} alkyl. It is preferred that R^5 and
15 R^6 both be hydrogen. It is also preferred that R^5 is hydrogen and R^6 is methyl. When R^5 is hydrogen and R^6 is methyl, both enantiomers and racemate are preferred.

"Alkyl" means a monovalent straight or branched chain group of
20 the formula C_xH_{2x+1} , with x being the number of carbon atoms.

"Alkenyl" means a straight or branched hydrocarbon radical containing a carbon-carbon double bond.

25 "Y — X" and "Y — Z" refer to a single bond or double bond attachment when defined by the substituents X, Y, and Z.

"Cycloalkyl" groups are monovalent cyclic moieties containing at least 3 carbon atoms and conforming to the formula $C_xH_{(2x-1)}$, with x
30 being the number of carbon atoms present. The cyclopropyl group is a preferred cycloalkyl moiety.

"Haloalkyl" includes straight and branched chain hydrocarbon radicals bearing from 1 to 3 halogen moieties. "Halogen" means F, Cl, Br or I. Preferred halogens in haloalkyl moieties of R⁴ include F and Cl.

- 5 One group of preferred compounds include the benzofurans and dihydrobenzofurans of Formula I wherein the group, -X-Y-Z-, consists of -CH₂-CH₂-O-, -CH=CH-O-, -CH₂-C(CH₃)₂-O-, -CH₂-CH(CH₃)-O- and -CH=CCH₃-O-.

- 10 Some preferred compounds of this group include:

(cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-propanamide;

- 15 (cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;

(cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-cyclopropane carboxamide;

- 20 (cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-butyramide; and

(cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-2-methyl propanamide.

25


Another group of preferred compounds include the benzodioxoles and benzodioxanes of Formula I wherein the group, -X-Y-Z-, consists of -O-CH₂-O- and -O-(CH₂)₂-O-, respectively.

- 30 Still another group of preferred compounds include the benzopyrans of Formula I wherein the group, -X-Y-Z-, consists of -CH₂-(CH₂)₂-O-.

Additionally, compounds of Formula I encompass all pharmaceutically acceptable solvates, particularly hydrates, thereof. The present invention also encompasses diastereomers as well as optical isomers, e.g. mixtures of enantiomers including racemic mixtures, as well as
5 individual enantiomers and diastereomers, which arise as a consequence of structural asymmetry in certain compounds of Formula I. Separation of the individual isomers or selective synthesis of the individual isomers is accomplished by application of various methods which are well known to practitioners in the art.

10

Compounds of Formula I can be prepared using the overall processes and many of several modifications shown in the following Reaction Scheme:

Reaction Scheme 1

- 5 The starting triflates of Formula 2 can be prepared by methods well known to those skilled in the art from the corresponding phenols. Conversion to the acetylenic alcohols of Formula 3 can be accomplished by palladium-mediated coupling to propargylic alcohol using a Pd catalyst like tetrakis(triphenylphosphine)palladium (0), palladium(II) acetate, dichlorobis(triphenylphosphine)palladium (II), and the like in the presence of a co-catalyst like copper iodide and a base like diisopropylamine in an inert solvent such as tetrahydrofuran or dimethylformamide. Reduction of the triple bond to the cis olefins of
- 10

Formula 4 can be accomplished by catalytic hydrogenation using palladium catalysts such as palladium on carbon in pyridine or Lindlar's catalyst. The cis olefin of Formula 4 can be converted to the cyclopropanes of Formula 5 using Simmons-Smith conditions such as treatment of diiodomethane with Zn-Cu couple or treatment of a dihalomethane with Et_2Zn in solvents such as ether or methylene chloride. The cyclopropyl alcohols of Formula 5 can then be converted to the amines of Formulas 8 and 10 by first forming the mesylates of Formula 6 with methane sulfonyl chloride and an acid scavenger, followed by displacement of the mesylate with sodium azide and reduction of the azides of Formula 7 with lithium aluminum hydride to provide the penultimate amines of Formula 8 or by displacement with sodium cyanide to give the compound of Formula 9, followed by reduction of the nitrile group with a reducing agent, such as LAH, to provide the penultimate amines of Formula 10. Further reaction of amines of Formulas 8 or 10 with acylating reagents provides compounds of Formula I. Suitable acylating agents include carboxylic acid halides, anhydrides, acyl imidazoles, alkyl isocyanates, alkyl isothiocyanates and carboxylic acids in the presence of condensing agents such as carbonyl imidazole, carbodiimides, and the like.

Biological Activity of the Compounds

The compounds of the invention are melatonergic agents. They have been found to bind human melatonergic receptors expressed in a stable cell line with good affinity. Further, the compounds are agonists as determined by their ability, like melatonin, to block the forskolin-stimulated accumulation of cAMP in certain cells. Due to these properties, the compounds and compositions of the invention should be useful as sedatives, chronobiotic agents, anxiolytics, antipsychotics, analgesics, and the like. Specifically, these agents should find use in the treatment of stress, sleep disorders, seasonal depression, appetite regulation, shifts in circadian cycles, melancholia, benign prostatic

hyperplasia, inflammatory articular diseases, headaches, and related conditions.

Melatonergic Receptor Binding Activity

5

1. Reagents:

- (a) TME = 50 mM Tris buffer containing 12.5 mM MgCl_2 , and 2 mM EDTA, pH 7.4 at 37°C with concentrated HCl.
- 10 (b) Wash buffer: 20 mM Tris base containing 2 mM MgCl_2 , pH 7.4 at room temperature.
- (c) 10^{-4} M melatonin (10^{-5} M final concentration).
- (d) 2-[^{125}I]-iodomelatonin, 0.1 M final concentration

15 2. Membrane Homogenates:

The melatonin ML_{1a} receptor cDNA was subcloned into pcDNA3 and introduced into NIH-3T3 cells using Lipofectamine. Transformed NIH-3T3 cells resistant to geneticin (G-418) were isolated, and single
20 colonies expressing high levels of 2-[^{125}I]-iodomelatonin binding were isolated. Cells are maintained in DMEM supplemented with 10% calf serum and G-418 (0.5 g/liter). Cells are grown to confluency in T-175 flasks, scraped using Hank's balanced salt solution, and frozen at -80°C. For preparing membrane homogenates, pellets are thawed on ice, and
25 resuspended in TME buffer in the presence of 10 $\mu\text{g/ml}$ aprotinin and leupeptin, and 100 μM phenylmethylsulfonylfluoride. The cells were then homogenized using a dounce homogenizer, and centrifuged. The resulting pellet was resuspended with dounce homogenizer in TME (supplemented with the above protease inhibitors) and frozen. On the
30 day of assay, the small aliquot was thawed on ice and resuspended in ice cold TME (1:50-1:100 v/v) and held on ice until assayed.

3. Incubation: 37°C for 1 hour. Reaction is terminated by filtration. Filters are washed 3 times.

4. References: Reppert, et al., Neuron, 13, p. 1177-1185 (1994).

5

Table 1

Example No.	Melatonin Binding
	Affinity (IC ₅₀) ^a
1	++
2	+
3	++
4	++
5	++

a = (IC₅₀) values for ML_{1a} human melatonin receptor binding

10

++ = <50 nM

+ = 50-200 nM

The compounds of the present invention have affinity for receptors of the endogenous pineal hormone, melatonin, as determined in a receptor binding assays described above in Table 1 for the ML_{1a} (human) receptors. Melatonin is involved in the regulation of a variety of biological rhythms and exerts its biological effects via interaction with specific receptors. There is evidence that administration of melatonin agonists are of clinical utility in the treatment of various conditions regulated by melatonin activity. Such conditions include depression, jet-lag, work-shift syndrome, sleep disorders, glaucoma, some disorders associated with reproduction, cancer, benign prostatic hyperplasia, immune disorders and neuroendocrine disorders.

For therapeutic use, the pharmacologically active compounds of Formula I will normally be administered as a pharmaceutical composition comprising as the (or an) essential active ingredient at least one such compound in association with a solid or liquid
5 pharmaceutically acceptable carrier and, optionally, with pharmaceutically acceptable adjuvants and excipients employing standard and conventional techniques.

The pharmaceutical compositions include suitable dosage forms
10 for oral, parenteral (including subcutaneous, intramuscular, intradermal and intravenous) transdermal, bronchial or nasal administration. Thus, if a solid carrier is used, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form, or in the form of a troche or lozenge. The solid carrier may contain conventional excipients such as
15 binding agents, fillers, tableting lubricants, disintegrants, wetting agents and the like. The tablet may, if desired, be film coated by conventional techniques. If a liquid carrier is employed, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule, sterile vehicle for injection, an aqueous or non-aqueous liquid suspension, or may be a
20 dry product for reconstitution with water or other suitable vehicle before use. Liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, wetting agents, non-aqueous vehicle (including edible oils), preservatives, as well as flavoring and/or coloring agents. For parenteral administration, a vehicle normally will
25 comprise sterile water, at least in large part, although saline solutions, glucose solutions and like may be utilized. Injectable suspensions also may be used, in which case conventional suspending agents may be employed. Conventional preservatives, buffering agents and the like also may be added to the parenteral dosage forms. Particularly useful is
30 the administration of a compound of Formula I in oral dosage formulations. The pharmaceutical compositions are prepared by conventional techniques appropriate to the desired preparation containing appropriate amounts of the active ingredient, that is, the compound of Formula I according to the invention. See, for example,

Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA, 17th edition, 1985.

In making pharmaceutical compositions containing compounds of the present invention, the active ingredient(s) will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, excipient or medium for the active ingredient.

Thus, the composition can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing for example up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders.

Some examples of suitable carriers and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, methyl cellulose, methyl- and propylhydroxybenzoates, talc, magnesium stearate and mineral oil. The formulations can additionally include lubricating agents, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The compositions of the invention may be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient.

The dosage of the compounds of Formula I to achieve a therapeutic effect will depend not only on such factors as the age, weight and sex of the patient and mode of administration, but also on the degree of melatonergic activity desired and the potency of the particular compound being utilized for the particular disorder or condition concerned. It is also contemplated that the treatment and dosage of the

particular compound may be administered in unit dosage form and that the unit dosage form would be adjusted accordingly by one skilled in the art to reflect the relative level of activity. The decision as to the particular dosage to be employed (and the number of times to be administered per day) is within the discretion of the physician, and may be varied by titration of the dosage to the particular circumstances of this invention to produce the desired therapeutic effect.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 0.1 to 100 mg, more usually 1 to 10 mg, of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier.

These active compounds are effective over a wide dosage range. For example, dosages per day will normally fall within the range of about 0.1 to 500 mg. In the treatment of adult humans, the range of about 0.1 to 10 mg/day, in single or divided doses, is preferred. Generally, the compounds of the invention may be used in treating sleep and related disorders in a manner similar to that used for melatonin.

However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances including the condition to be treated, the choice of compound to be administered, the chosen route of administration, the age, weight, and response of the individual patient, and the severity of the patient's symptoms.

The compounds which constitute this invention, their methods of preparation and their biologic actions will appear more fully from consideration of the following examples, which are given for the purpose

of illustration only and are not to be construed as limiting the invention in sphere or scope.

5 DESCRIPTION OF SPECIFIC EMBODIMENTS

In the following examples, used to illustrate the foregoing synthetic processes, all temperatures are expressed in degrees Celsius and melting points are uncorrected. Proton magnetic resonance (1H NMR) and carbon magnetic resonance (13C NMR) spectra were determined in the solvents indicated and chemical shifts are reported in δ units downfield from the internal standard tetramethylsilane (TMS) and interproton coupling constants are reported in Hertz (Hz). Splitting patterns are designated as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad peak; dd, doublet of doublet; bd, broad doublet; dt, doublet of triplet; bs, broad singlet; dq, doublet of quartet. The infrared (IR) spectral descriptions include only absorption wave numbers (cm⁻¹) having functional group identification value. The IR determinations were employed using the compound neat as a film or by employing potassium bromide (KBr) as diluent. Optical rotations $[\alpha]_D^{25}$ were determined in the solvents and concentration indicated. Low resolution mass spectra (MS) are reported as the apparent molecular weight (M+H)⁺. The elemental analyses are reported as percent by weight.


25

Preparation of Intermediates of Formula 2

(E)-N-[3-(2,3-Dihydrobenzofuran-4-yl)prop-2-enyl]-propanamide


30

Step 1: 2-Bromoresorcinol



Bromine (0.363 L) was added dropwise over 2 hours to a solution of resorcinol (250 g) in dichloromethane (3.5 L). The solution was stirred at room temperature for 18 hours at which time approximately 1 L of the dichloromethane was removed by distillation. MeOH was added and the distillation continued in this manner until all of the dichloromethane was removed and the solution contained approximately 1.5 L of MeOH. To this was added a solution of NaOH (181.5 g) and Na₂SO₃ (573 g) in H₂O (7.5 L). The resulting mixture was stirred at ambient temperature for 1 hour. The solution was then acidified to pH = 2 with concentrated HCl (75 mL) and extracted with tert-butyl methyl ether (TBME) (2 X 1L). The combined organic layers were treated with activated charcoal (20 g) and filtered through Celite; the Celite was washed with an additional 500 mL of TBME. The solvent was then removed in vacuo. The resulting crude 2-bromoresorcinol was dissolved in a minimum amount of ethyl acetate and filtered through silica gel eluting with a gradient from 20% to 40% ethyl acetate in hexanes yielding 2-bromoresorcinol (122 g). m.p. 86-88°.

Step 2: 2,6-Di(2-chloroethoxy)bromobenzene




Potassium carbonate (536 g), sodium iodide (9.76 g), and sodium metabisulfate (12.2 g) were suspended in dichloromethane (1.53 L) and DMF (0.4 L) and heated to 80°C. A solution of 2-bromoresorcinol (122.3 g) in DMF (0.4 L) was then added dropwise over 2 hours. The reaction was stirred at 80°C for 20 hours, cooled to ambient temperature and

filtered through a medium porosity fritted funnel. The solid residue was washed with DMF (2 X 0.28 L) and organic fractions combined. The organics were washed with 1N HCl (1 X 1.84 L and 1 X 0.92 L), half-saturated NaHCO₃ solution (0.92 L), half-saturated brine (0.92 L), dried
5 over Na₂SO₄ and concentrated in vacuo.

The crude product was dissolved in EtOH (142 mL) and TBME (76 mL) and treated with activated charcoal (14 g) at 70°C for 0.25 hours. The suspension was filtered through Celite. The solution was cooled to
10 0°C for 48 hours and the crystals collected (44.88 g). The mother liquor was concentrated and passed over a plug of silica gel eluting with 20% ethyl acetate in hexane yielding an additional 39.25 g of pure product. The total yield of title compound was 84.13 g, (41.4%).

15 **Step 3:** 2,3-Dihydro-4-hydroxybenzofuran



The product of Step 2 (39.25 g) was dissolved in THF (0.5 L) and
20 cooled to -78°C. nBuLi (2.5M in hexanes, 300 mL) was then added dropwise over 30 minutes and the reaction stirred at -70°C for an additional 45 minutes. The solution was then warmed to 0°C over 10 minutes and stirred at this temperature for 1 hour. Glacial acetic acid (16 mL) was added followed by 1N NaOH (160 mL) and the layers allowed
25 to separate. The organics were extracted with 1N NaOH (2 X 80 mL) and the combined aqueous fractions were then washed with TBME (160 mL). TBME (240 mL) was then added and the aqueous layer acidified with 6N HCl. The aqueous layer was re-extracted with TBME (240 mL), and the combined organics stirred over activated charcoal (5 g) for 15
30 minutes, filtered through Celite, and concentrated in vacuo. The crude

product was crystallized from toluene and heptane to yield the title compound (16.8 g, 99%).

Step 4: 2,3-Dihydrobenzofuran-4-yl trifluoromethane sulfonate

5



The product of Step 3 (1.0 g) was dissolved in anhydrous dichloromethane (5 mL) and cooled to 0°C. Pyridine (0.87 mL) was then added followed by dropwise addition over 30 minutes of trifluoromethansulfonic anhydride (2.28 g). Stirred from 0°C to ambient temperature over 1 hour. The methylene chloride solution was then washed with water (2 X 4.6 mL), 10% phosphoric acid (4.6 mL), saturated NaHCO₃ solution (4.6 mL) and brine (2.3 mL). The solution was then treated with activated carbon (170 mg) for 5 minutes, filtered through Celite, dried over Na₂SO₄ and concentrated in vacuo to yield the title compound (1.74g, 88%).

Preparation of Intermediates of Formula 3

20

3-(2,3-Dihydrobenzofuran-4-yl)prop-2-yn-1-ol

A mixture of the triflate (0.1 mol), propargylic alcohol (0.2 mol), CuI (10 mmol), dichlorobis(triphenylphosphine)palladium (II) (5 mmol) and diisopropylamine (100 mL) in 300 mL of DMF was stirred at 80°C for 15 h and then at RT for 4 days. The crude reaction mixture was then diluted with ether (300 mL) and washed sequentially with water (1 x 200 mL), 1N HCl (3 x 200 mL), saturated aqueous sodium bicarbonate solution (1 x 200 mL) and then brine (1 x 200 mL). The organic phase was separated, dried over magnesium sulfate, filtered and concentrated in vacuo to provide a brown oil. Purification by silica gel column

chromatography (Hexanes:EtOAc, 4:1) provided 6.44 g (37%) of the desired product as a yellow-brown oil.

¹H NMR (300 MHz, CDCl₃) δ 7.07 (t, J=7.8 Hz, 1H), 6.93 (d, J=7.6 Hz, 1H), 6.77 (d, J=7.9 Hz, 1H), 4.59 (t, J=8.8 Hz, 2H), 4.52 (s, 2H), 3.26 (t, J=8.7 Hz, 2H).

Preparation of Intermediates of Formula 4

10 (cis)-3-(2,3-Dihydrobenzofuran-4-yl)prop-2-en-1-ol

A mixture of 3-(2,3-dihydrobenzofuran-4-yl)prop-2-yn-1-ol (6.2 mmol) and Pd/C (300 mg) in 20 mL of pyridine was shaken under an atmosphere of H₂ (10 psi) until no starting material remained (8 h). The crude reaction mixture was filtered to remove the catalyst and the catalyst was washed with ether. The filtrate was concentrated in vacuo to provide 1.1 g of the desired product as a yellow oil (quantitative yield).

¹H NMR (300 MHz, CDCl₃) δ 7.11 (t, J=7.8 Hz, 1H), 6.74 (d, J=8.0 Hz, 1H), 6.68 (d, J=7.7 Hz, 1H), 6.48 (d, J=11.7 Hz, 1H), 5.89-5.97 (m, 1H), 4.59 (t, J=8.7 Hz, 2H), 4.42 (dd, J=6.4, 1.6 Hz, 2H), 3.15 (t, J=8.7 Hz, 2H).

Preparation of Intermediates of Formula 5

25 (cis)-2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl methanol

Diethylzinc (63 mmol, 1.0 M in hexanes) was added dropwise to a mixture of (cis)-3-(2,3-dihydrobenzofuran-4-yl)prop-2-en-1-ol (12.6 mmol) in 10 mL of CH₂Cl₂ at -15°C. The resulting mixture was stirred at -15°C for 1 h followed by the dropwise addition of CH₂I₂ (63 mmol). The resulting mixture was warmed to RT and stirred overnight. The crude reaction mixture was then quenched with saturated aqueous NH₄Cl

organic extracts were washed with saturated aqueous NH_4Cl solution, dried over magnesium sulfate, filtered and concentrated in vacuo. Purification by silica gel column chromatography (Hexanes:EtOAc, 4:1) provided 2.06 g (86%) of the desired product.

5

^1H NMR (300 MHz, CDCl_3) δ 7.05 (t, $J=7.9$ Hz, 1H), 6.68 (d, $J=7.9$ Hz, 1H), 6.61 (d, $J=7.7$ Hz, 1H), 4.57-4.64 (m, 2H), 3.43-3.49 (m, 1H), 3.24-3.35 (m, 3H), 2.06-2.15 (m, 1H), 1.52-1.59 (m, 1H), 1.02-1.10 (m, 1H), 0.89-0.94 (m, 1H).

10

Preparation of Intermediates of Formula 6

(cis)-[2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl methane sulfonate

15

Methane sulfonyl chloride (6.5 mmol) was added dropwise to a solution of (cis)-2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl methanol (5.0 mmol) and triethylamine (7.5 mmol) in 50 mL of CH_2Cl_2 at 0°C . The resulting mixture was allowed to warm to RT and stirred for 2 h. The crude reaction mixture was then diluted with CH_2Cl_2 (50 mL) and washed sequentially with water (2 x 50 mL), saturated aqueous sodium bicarbonate solution (2 x 50 mL) and then brine (1 x 50 mL). The organic phase was separated, dried over magnesium sulfate, filtered and concentrated in vacuo to provide 1.38 g of the desired product as an oil (quantitative yield).

20

^1H NMR (300 MHz, CDCl_3) δ 6.98 (t, $J=7.8$ Hz, 1H), 6.61 (d, $J=7.9$ Hz, 1H), 6.49 (d, $J=7.7$ Hz, 1H), 4.51-4.57 (m, 2H), 4.11 (dd, $J=6.1, 10.5$ Hz, 1H), 3.52-3.58 (m, 1H), 3.15-3.21 (m, 2H), 2.48 (s, 3H), 2.13-2.19 (m, 1H), 1.57-1.69 (m, 1H), 1.07-1.15 (m, 1H), 0.94-1.00 (m, 1H).

30

Preparation of Intermediates of Formula 7

(cis)-2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl methyl azide

A solution of (cis)-[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl methanesulfonate (5.0 mmol) and sodium azide (10.0 mmol) in 100 mL of dimethylformamide was heated at 70°C for 1 h. The reaction mixture was allowed to cool to room temperature and stirred overnight. The crude reaction mixture was then diluted with CH₂Cl₂ (100 mL) and washed sequentially with water (3 x 100 mL) and then brine (1 x 100 mL). The organic phase was separated, dried over magnesium sulfate, filtered and concentrated in vacuo to provide 1.1 g of the desired product as an oil (quantitative yield).

¹H NMR (300 MHz, CDCl₃) δ 7.07 (t, J=7.8 Hz, 1H), 6.70 (d, J=7.9 Hz, 1H), 6.56 (d, J=7.6 Hz, 1H), 4.59-4.65 (m, 2H), 3.24-3.30 (m, 2H), 3.07 (dd, J=6.9, 13 Hz, 1H), 2.95 (dd, J=7.9, 13 Hz, 1H), 2.12-2.19 (m, 1H), 1.51-1.57 (m, 1H), 1.12-1.19 (m, 1H), 0.93-0.99 (m, 1H).

Preparation of Intermediates of Formula 8

(cis)-2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl methyl amine

A solution of (cis)-2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl methyl azide (5.0 mmol) in 20 mL of tetrahydrofuran was added dropwise to a mixture of lithium aluminum hydride (10.0 mmol) in 40 mL of tetrahydrofuran at -40°C. The resulting mixture was allowed to warm to RT and stirred overnight. The crude reaction mixture was quenched by the sequential addition of water (1.0 mL), 10 N sodium hydroxide (1.0 mL) and water (3.0 mL). The solid material was removed by filtration through Celite and the filtrate was concentrated in vacuo. The resulting residue was dissolved in ether (100 mL) and washed with brine (50 mL). The organic phase was separated, dried over magnesium sulfate,

filtered and concentrated in vacuo to provide 630 mg of the desired product as an oil (66%).

¹H NMR (300 MHz, CDCl₃) δ 7.04 (t, J=7.8 Hz, 1H), 6.67 (d, J=7.9 Hz, 1H), 6.57 (d, J=7.7 Hz, 1H), 4.55-4.63 (m, 2H), 3.22-3.29 (m, 2H), 2.38-2.52 (m, 2H), 2.01-2.07 (m, 1H), 1.35-1.45 (m, 3H), 0.98-1.05 (m, 1H), 0.81-0.87 (m, 1H).

EXAMPLE 1

10

(cis)-N-[[2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide

A solution of (cis)-2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl methyl amine (0.43 mmol), triethylamine (1.30 mmol) and propionyl chloride (0.65 mmol) in 10 mL of methylene chloride was stirred at room temperature overnight. The reaction mixture was washed with brine (2 x 20 mL). The organic phase was separated, dried over magnesium sulfate, filtered and concentrated in vacuo. Purification by silica gel column chromatography (Hex:EtOAc, 4:1) provided 23 mg of the desired product (22%) as a yellow oil. HPLC purity > 95%.

¹H NMR (300 MHz, CDCl₃) δ 7.04 (t, J=7.8 Hz, 1H), 6.67 (d, J=7.9 Hz, 1H), 6.57 (d, J=7.6 Hz, 1H), 5.27 (br s, 1H), 4.56-4.64 (m, 2H), 3.12-3.33 (m, 2H), 2.98-3.04 (m, 2H), 2.12 (q, J=7.6 Hz, 2H), 2.00-2.18 (m, 1H), 1.42-1.55 (m, 1H), 1.09 (t, J=7.6 Hz, 3H), 0.99-1.21 (m, 1H), 0.86-0.92 (m, 1H);

¹³C NMR (75 MHz, CDCl₃) δ 173.7, 160.0, 135.4, 128.2, 127.4, 119.5, 107.8, 71.2, 39.7, 29.9, 29.2, 18.8, 18.0, 10.0, 8.5;

IR (neat) 1643, 1548, 1235 cm⁻¹;

MS (ESI) *m/e* 246 (M+H)⁺.

EXAMPLE 2**(cis)-N-[[2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide**

5

The title compound was prepared by the general procedure described in Example 1 using (cis)-2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl methyl amine (0.79 mmol), triethylamine (2.37 mmol) and acetyl chloride (0.95 mmol). Purification by silica gel column chromatography (Hex:EtOAc, 1:1) provided 92 mg of the desired product (50%) as an off-white wax. HPLC purity = 93%.

¹H NMR (300 MHz, CDCl₃) δ 6.97 (t, J=7.8 Hz, 1H), 6.59 (d, J=7.9 Hz, 1H), 6.49 (d, J=7.7 Hz, 1H), 5.28 (br s, 1H), 4.46-4.58 (m, 2H), 2.94-3.25 (m, 3H), 2.79-2.88 (m, 1H), 1.94-2.04 (m, 1H), 1.83 (s, 3H), 1.34-1.47 (m, 1H), 0.92-1.00 (m, 1H), 0.77-0.81 (m, 1H);

¹³C NMR (75 MHz, CDCl₃) δ 170.0, 159.9, 135.2, 128.1, 127.3, 119.5, 107.7, 71.1, 39.8, 29.0, 23.4, 18.7, 17.8, 8.4;

IR (KBr) 3329, 1648, 1557, 1236 cm⁻¹;

MS (ESI) *m/e* 232 (M+H)⁺.

EXAMPLE 3**(cis)-N-[[2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide**

The title compound was prepared by the general procedure described in Example 1 using (cis)-2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl methyl amine (0.79 mmol), triethylamine (2.37 mmol) and cyclopropane carbonyl chloride (0.95 mmol). Purification by silica gel column chromatography (Hex:EtOAc, 2:1) provided 88 mg of the desired product (43%) as a white wax. HPLC purity = 92%.

¹H NMR (300 MHz, CDCl₃) δ 7.02 (t, J=7.8 Hz, 1H), 6.65 (d, J=8.0 Hz, 1H), 6.56 (d, J=7.7 Hz, 1H), 5.45 (br s, 1H), 4.45-4.62 (m, 2H), 3.12-3.33 (m, 2H), 2.97-3.04 (m, 2H), 1.99-2.07 (m, 1H), 1.44-1.54 (m, 1H), 1.15-1.21 (m, 1H), 0.98-1.05 (m, 1H), 0.85-0.92 (m, 3H), 0.64-0.70 (m, 2H);

5

¹³C NMR (75 MHz, CDCl₃) δ 173.4, 159.8, 135.3, 128.1, 127.4, 119.4, 107.6, 71.1, 39.9, 29.1, 18.7, 17.9, 14.8, 8.4, 7.1;

IR (KBr) 3323, 1639, 1558, 1238 cm⁻¹;

MS (ESI) *m/e* 258 (M+H)⁺.

10

EXAMPLE 4

15 (cis)-N-[[2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butyramide

The title compound was prepared by the general procedure described in Example 1 using (cis)-2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl methyl amine (0.79 mmol), triethylamine (2.37 mmol) and butyryl chloride (0.95 mmol). Purification by silica gel column chromatography (Hex:EtOAc, 2:1) provided 61 mg of the desired product (30%) as a white solid. HPLC purity = 98%. m.p. 73-74°;

25 ¹H NMR (300 MHz, CDCl₃) δ 6.97 (t, J=7.8 Hz, 1H), 6.59 (d, J=8.0 Hz, 1H), 6.50 (d, J=7.7 Hz, 1H), 5.19 (br s, 1H), 4.46-4.59 (m, 2H), 3.02-3.26 (m, 2H), 2.85-3.02 (m, 2H), 1.94-2.01 (m, 3H), 1.46-1.61 (m, 2H), 1.36-1.46 (m, 1H), 0.90-0.99 (m, 1H), 0.84 (t, J=7.3 Hz, 3H), 0.78-0.87 (m, 1H);

30 ¹³C NMR (75 MHz, CDCl₃) δ 172.8, 159.9, 135.3, 128.1, 127.3, 119.4, 107.6, 71.1, 39.6, 38.8, 29.0, 19.2, 18.7, 17.9, 13.8, 8.4;

IR (KBr) 3312, 1640, 1554, 1237 cm⁻¹;

MS (ESI) *m/e* 260 (M+H)⁺.

EXAMPLE 5**(cis)-N-[[2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-2-methyl propanamide**

5

The title compound was prepared by the general procedure described in Example 1 using (cis)-2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl methyl amine (0.79 mmol), triethylamine (2.37 mmol) and isobutyryl chloride (0.95 mmol). Purification by recrystallization from
10 EtOAc/Hexanes provided 76 mg of the desired product (37%) as colorless plates. HPLC purity > 98%. m.p. 125-126°;

¹H NMR (300 MHz, CDCl₃) δ 6.97 (t, J=7.8 Hz, 1H), 6.59 (d, J=7.9 Hz, 1H), 6.50 (d, J=7.7 Hz, 1H), 5.17 (br s, 1H), 4.46-4.59 (m, 2H), 3.04-3.26
15 (m, 2H), 2.92-2.97 (m, 2H), 2.16 (septet, J=6.9 Hz, 1H), 1.94-2.01 (m, 1H), 1.34-1.47 (m, 1H), 1.01 (dd, J=2.2, 6.9 Hz, 6H), 0.93-1.06 (m, 1H), 0.80-0.85 (m, 1H);

¹³C NMR (75 MHz, CDCl₃) δ 176.7, 159.9, 135.3, 128.1, 127.2, 119.3,
20 107.6, 71.1, 39.5, 35.7, 29.1, 19.6, 18.6, 17.9, 8.4;


IR (KBr) 3309, 1640, 1553, 1238 cm⁻¹;

MS (ESI) *m/e* 260 (M+H)⁺.

We claim:

1. A cis cyclopropane compound of Formula I or a pharmaceutically acceptable solvate thereof having the formula:

5



wherein

- R¹ and R² each are independently hydrogen or halogen;
 X is CH₂, CH or oxygen;
 10 Y is CR⁵, CR⁵R⁶ or (CH₂)_n, with n = 1-2;
 Z is CH₂, CH or oxygen;
 m is 1 or 2;
 R³ is hydrogen or C₁₋₄ alkyl;
 R⁴ is C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₃ haloalkyl, C₂₋₆ alkenyl,
 15 C₁₋₄ alkoxy(C₁₋₄)alkyl, C₁₋₄ alkylthio(C₁₋₄)alkyl or C₁₋₄ trifluoromethylalkyl; and
 R⁵ and R⁶ each are independently hydrogen or C₁₋₄ alkyl.

2. A compound of claim 1 wherein R³ is hydrogen and m is 1.
 20 3. A compound of claim 2 wherein R⁴ is C₁₋₆ alkyl or C₃₋₄ cycloalkyl.
 4. A compound of claim 3 wherein X is CH₂ and Z is oxygen.
 25 5. A compound of claim 4 selected from the group consisting of:

(cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-propanamide;

(cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;

(cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-
cyclopropane carboxamide;

5

(cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-
butyramide; and

10 (cis)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-2-methyl
propanamide.

6. The compound of claim 5 which is (cis)-N-[[2-(2,3-dihydrobenzo-
furan-4-yl)cycloprop-1-yl]methyl]propanamide.

15 7. A method of treating sleep disorders in a patient in need of such
treatment comprising administering to said patient a therapeutic amount
of a compound of claim 1.

20 8. A composition useful for treating sleep disorders comprising a
therapeutic amount of a compound of claim 1 and a suitable amount of a
pharmaceutically acceptable carrier.

25 9. A method of treating circadian rhythm-related disorders in a
patient in need of such treatment comprising administering to said
patient a therapeutic amount of a compound of claim 1.

10. A composition useful for treating circadian rhythm-related
disorders comprising a therapeutic amount of a compound of claim 1
and a suitable amount of a pharmaceutically acceptable carrier.

30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/11474

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 514/452, 464, 469; 549/365, 441, 462

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/452, 464, 469; 549/365, 441, 462

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 98/25606 A1 (BRISTOL-MYERS SQUIBB COMPANY) 18 June 1998, see the entire document especially col. 9, lines 5-17.	1-10
X,P	US 5,856,529A [CATT et al.] 05 January, 1999, see the entire document especially col. 5, lines 30-50.	1-10

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A document defining the general state of the art which is not considered to be of particular relevance	*X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B earlier document published on or after the international filing date	*Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A document member of the same patent family
*O document referring to an oral disclosure, use, exhibition or other means	
*P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

21 JULY 1999

Date of mailing of the international search report

21 OCT 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

TAOFIQ A. SOLOTA

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/11474

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (6):

A61K 31/34, 31/36, 31/335; A01N 43/08, 43/30, 43/32; C07D 307/88, 317/46, 319/04, 319/08